



# Cambridge International AS & A Level

---

## MATHEMATICS

9709/11

Paper 1 Pure Mathematics 1

May/June 2026

MARK SCHEME

Maximum Mark: 75

---

Published

This mark scheme contains the instructions and marking guidance that our examiners used to mark candidate responses for this component.

Before they start marking, examiners complete a standardisation process. They review a range of candidate responses and discuss the mark scheme in detail to agree which answers can and cannot be accepted. Examiners award marks where it is clear that candidate responses have met the requirements of the mark scheme. These requirements may vary between different components or different versions of the same component, depending on the exact requirements of the question and the candidate responses seen during the standardisation process.

Sometimes, our mark schemes may allow marks to be awarded to responses which contain elements that are factually incorrect or for content not covered by the syllabus. We do this in response to the demands of a specific question to support candidates and make sure that our assessments are fair. However, these allowances should not be used as a guide for teaching and learning.

We cannot discuss the mark scheme with schools, and we recommend that schools read the mark scheme together with the assessment material and the Principal Examiner Report for Teachers.

---

This document consists of **21** printed pages.

**General marking principles**

All examiners must apply these marking principles when marking candidate responses.

|   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|---|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1 | Examiners must award marks for each response in line with: <ul style="list-style-type: none"> <li>the specific <b>content</b> defined in the mark scheme</li> <li>the specific <b>skills</b> defined in the mark scheme or in the level descriptions</li> <li>the <b>standard of response</b> required as exemplified by the standardisation scripts.</li> </ul>                                                                                                                                                                                                                                                                                    |
| 2 | Examiners must award <b>whole marks</b> (not half marks, or other fractions).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| 3 | Examiners must award marks <b>positively</b> . This means that: <ul style="list-style-type: none"> <li>marks are not deducted for errors or omissions</li> <li>marks are awarded for correct/valid answers as defined in the mark scheme and also for <b>other valid answers</b> which go beyond the scope of the syllabus and mark scheme referring to your team leader as appropriate</li> <li>the quality of spelling, punctuation and grammar are judged <b>only</b> when the mark scheme indicates that these features are <b>specifically assessed</b> in the question. However, the meaning of all responses must be unambiguous.</li> </ul> |
| 4 | Examiners must <b>consistently</b> apply the requirements of the mark scheme and any rules for what to do when candidates have not followed instructions correctly.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| 5 | Examiners must use the <b>full range of marks</b> defined in the mark scheme, unless limited by the quality of the candidate responses seen.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| 6 | Examiners must award marks according to the mark scheme and <b>must not</b> award marks with grade thresholds or grade descriptions in mind.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |

**Mathematics-Specific Marking Principles**

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- 5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

**Annotations guidance for centres**

Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standardisation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.

We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

**Annotations**

| <b>Annotation</b>                                                                   | <b>Meaning</b>                         |
|-------------------------------------------------------------------------------------|----------------------------------------|
|    | More information required              |
|    | Accuracy mark awarded zero             |
|    | Accuracy mark awarded one              |
|    | Independent accuracy mark awarded zero |
|  | Independent accuracy mark awarded one  |
|  | Independent accuracy mark awarded two  |
|  | Benefit of the doubt                   |
|  | Blank Page                             |
|  | Incorrect                              |
| Dep                                                                                 | Used to indicate DM0 or DM1            |

| <b>Annotation</b>                                                                   | <b>Meaning</b>                                                                                                                             |
|-------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|
| DM1                                                                                 | Dependent on the previous M1 mark(s)                                                                                                       |
|    | Follow through                                                                                                                             |
|    | Indicate working that is right or wrong                                                                                                    |
| Highlighter                                                                         | Highlight a key point in the working                                                                                                       |
|    | Ignore subsequent work                                                                                                                     |
|    | Judgement                                                                                                                                  |
|    | Judgement                                                                                                                                  |
|    | Method mark awarded zero                                                                                                                   |
|    | Method mark awarded one                                                                                                                    |
|    | Method mark awarded two                                                                                                                    |
|    | Misread                                                                                                                                    |
|    | Omission or Other solution                                                                                                                 |
| Off-page comment                                                                    | Allows comments to be entered at the bottom of the RM marking window and then displayed when the associated question item is navigated to. |
| On-page comment                                                                     | Allows comments to be entered in speech bubbles on the candidate response.                                                                 |
|  | Judgment made by the PE                                                                                                                    |
|  | Premature approximation                                                                                                                    |
|  | Special case                                                                                                                               |
|  | Indicates that work/page has been seen                                                                                                     |

| Annotation                                                                        | Meaning                                |
|-----------------------------------------------------------------------------------|----------------------------------------|
|  | Error in number of significant figures |
|  | Correct                                |
|  | Transcription error                    |
|  | Correct answer from incorrect working  |

**PUBLISHED****Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

**Types of mark**

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
  - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
  - The total number of marks available for each question is shown at the bottom of the Marks column.
  - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
  - Square brackets [ ] around text or numbers show extra information not needed for the mark to be awarded.

**Abbreviations**

|        |                                                                                                                                                                                       |
|--------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| AEF/OE | Any Equivalent Form (of answer is equally acceptable) / Or Equivalent                                                                                                                 |
| AG     | Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)                                                   |
| CAO    | Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)                                                                                           |
| CWO    | Correct Working Only                                                                                                                                                                  |
| ISW    | Ignore Subsequent Working                                                                                                                                                             |
| SOI    | Seen Or Implied                                                                                                                                                                       |
| SC     | Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance) |
| WWW    | Without Wrong Working                                                                                                                                                                 |
| AWRT   | Answer Which Rounds To                                                                                                                                                                |

## PUBLISHED

| Question | Answer                                                                                                                | Marks | Guidance                                                                                  |
|----------|-----------------------------------------------------------------------------------------------------------------------|-------|-------------------------------------------------------------------------------------------|
| 1        | <b>EITHER</b><br>Solution 1<br>$r^2 = \frac{12.5}{50} \quad [\Rightarrow r = \frac{1}{2}]$                            | (M1)  | Correct procedure to find $r^2$ or $r$ .                                                  |
|          | $a = 200$ and $r = \frac{1}{2}$                                                                                       | A1    | Ignore $r = -\frac{1}{2}$ .                                                               |
|          | $S_{\infty} = \frac{200}{1 - \frac{1}{2}}$                                                                            | M1    | Use of formula for sum to infinity for <i>their</i> $a$ and $r$ provided that $ r  < 1$ . |
|          | $S_{\infty} = 400$                                                                                                    | A1)   | CAO                                                                                       |
|          | <b>OR</b><br>Solution 2<br><b>Alternative method for M1 A1</b><br>$a = \frac{2500}{12.5} \quad [\Rightarrow a = 200]$ | (M1)  | Correct procedure to find $a$ .                                                           |
|          | $r = \frac{1}{2}$ and $a = 200$                                                                                       | A1)   |                                                                                           |
|          | Available marks                                                                                                       | 4     |                                                                                           |

## PUBLISHED

| Question | Answer                                                  | Marks       | Guidance                                                                                                    |
|----------|---------------------------------------------------------|-------------|-------------------------------------------------------------------------------------------------------------|
| 2(a)(i)  | $(1+2x)^5 = 1+10x+40x^2$                                | <b>B1</b>   | B1 for any two.                                                                                             |
|          |                                                         | <b>B1</b>   | B1 for all three.<br>ISW for extra terms.                                                                   |
|          |                                                         | <b>2</b>    |                                                                                                             |
| 2(a)(ii) | $(3+x^2)^6 = 729+1458x^2$                               | <b>B1</b>   | ISW extra terms, e.g. $1215x^4$ .                                                                           |
|          |                                                         | <b>1</b>    |                                                                                                             |
| 2(b)     | Coefficient of $x^2$ is $1 \times 1458 + 40 \times 729$ | <b>M1</b>   | For using <i>their</i> values to find coefficient<br>from $(ax^2 + bx + c)(d + ex^2) \Rightarrow ad + ce$ . |
|          | 30618                                                   | <b>A1FT</b> | Condone $30618x^2$ .                                                                                        |
|          |                                                         | <b>2</b>    |                                                                                                             |

| Question | Answer                                 | Marks     | Guidance                                                            |
|----------|----------------------------------------|-----------|---------------------------------------------------------------------|
| 3        | $y = \left(\frac{x-5}{4}\right)^3 - 2$ | <b>B1</b> | B1 for $x-5$ , or subtracting 2 or dividing by 4 <sup>3</sup> .     |
|          |                                        | <b>B1</b> | Second B1 for another correct procedure.                            |
|          |                                        | <b>B1</b> | Final B1 for fully correct OE, e.g. $y = \frac{1}{64}(x-5)^3 - 2$ . |
|          |                                        | <b>3</b>  |                                                                     |

## PUBLISHED

| Question | Answer                                                                                                                                                                            | Marks        | Guidance                                                                                                                                       |
|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|------------------------------------------------------------------------------------------------------------------------------------------------|
| 4        | <b>EITHER</b><br>Solution 1<br>$8\sin^4\theta - 6(1 - \sin^2\theta) + 1 = 0 \quad [\Rightarrow 8\sin^4\theta + 6\sin^2\theta - 5 = 0]$                                            | <b>(M1)*</b> | Use of $\sin^2\theta + \cos^2\theta = 1$ to eliminate $\cos^2\theta$ .                                                                         |
|          | $(4\sin^2\theta + 5)(2\sin^2\theta - 1) = 0 \quad [\Rightarrow \sin^2\theta = \frac{1}{2} \text{ or } \sin^2\theta = -\frac{5}{4}]$                                               | <b>DM1</b>   | Solve <i>their</i> three-term quadratic in $\sin^2\theta$ using suitable method, e.g. factorisation, quadratic formula, completing the square. |
|          | $\sin\theta = \pm \frac{1}{\sqrt{2}}$                                                                                                                                             | <b>A1</b>    | <b>SC B1</b> if M1 M0, allow AWRT $\pm 0.707$ .                                                                                                |
|          | $\theta = \frac{1}{4}\pi, \frac{3}{4}\pi, \frac{5}{4}\pi, \frac{7}{4}\pi$ or 0.785, 2.36, 3.93, 5.50                                                                              | <b>A1</b>    | A1 for any correct pair.                                                                                                                       |
|          |                                                                                                                                                                                   | <b>A1)</b>   | A1 for all four and no other solutions in range.<br><b>SC B1</b> for 45°, 135°, 225°, 315°.                                                    |
|          | <b>OR</b><br>Solution 2<br><b>Alternative Method for Question 4</b><br>$8(1 - \cos^2\theta)^2 - 6\cos^2\theta + 1 = 0 \quad [\Rightarrow 8\cos^4\theta - 22\cos^2\theta + 9 = 0]$ | <b>(M1)*</b> | Use of $\sin^2\theta + \cos^2\theta = 1$ to eliminate $\sin^2\theta$ .                                                                         |
|          | $(4\cos^2\theta - 9)(2\cos^2\theta - 1) = 0 \quad [\Rightarrow \cos^2\theta = \frac{9}{4} \text{ or } \cos^2\theta = \frac{1}{2}]$                                                | <b>DM1</b>   | Solve three-term quadratic in $\cos^2\theta$ using suitable method, e.g. factorisation, quadratic formula, completing the square.              |
|          | $\cos\theta = \pm \frac{1}{\sqrt{2}}$                                                                                                                                             | <b>A1</b>    | <b>SC B1</b> if M1 M0, allow AWRT $\pm 0.707$ .                                                                                                |
|          | $\theta = \frac{1}{4}\pi, \frac{3}{4}\pi, \frac{5}{4}\pi, \frac{7}{4}\pi$                                                                                                         | <b>A1</b>    | A1 for any correct pair.                                                                                                                       |
|          |                                                                                                                                                                                   | <b>A1)</b>   | A1 for all four and no other solutions in range.<br><b>SC B1</b> for 45°, 135°, 225°, 315°.                                                    |
|          | Available marks                                                                                                                                                                   | <b>5</b>     |                                                                                                                                                |

## PUBLISHED

| Question | Answer                                                                                                                            | Marks       | Guidance                                                                            |
|----------|-----------------------------------------------------------------------------------------------------------------------------------|-------------|-------------------------------------------------------------------------------------|
| 5        | $\left[\frac{dy}{dx}\right] = -0.5 + \frac{1}{2}(x^3 - 3x - 2)^{-\frac{1}{2}}(3x^2 - 3)$                                          | <b>B1B1</b> | B1 for $\frac{1}{2}(x^3 - 3x - 2)^{-\frac{1}{2}}$ , B1 for the rest.                |
|          | At $x = 6$ , $\left[\frac{dy}{dx}\right] = -0.5 + \frac{1}{2} \times 196^{-\frac{1}{2}} \times 105 \left[ = \frac{13}{4} \right]$ | <b>M1</b>   | For attempt at gradient by substituting $x = 6$ into <i>their</i> $\frac{dy}{dx}$ . |
|          | [Gradient of normal] = $-\frac{4}{13}$                                                                                            | <b>M1</b>   | For use of product of gradients = $-1$ using <i>their</i> $\frac{dy}{dx}$ .         |
|          | Equation is $4x + 13y - 167 = 0$                                                                                                  | <b>A1</b>   | OE in required form.                                                                |
|          |                                                                                                                                   | <b>5</b>    |                                                                                     |

## PUBLISHED

| Question | Answer                                                                                                                                | Marks        | Guidance                                                                                   |
|----------|---------------------------------------------------------------------------------------------------------------------------------------|--------------|--------------------------------------------------------------------------------------------|
| 6        | <b>EITHER</b><br>Solution 1<br>$p^2 + (p+6) = 2 \times 4p$ or $4p - (p+6) = p^2 - 4p$<br>$[\Rightarrow p^2 - 7p + 6 = 0]$             | <b>(M1)*</b> | OE<br>Correct method for forming quadratic equation.<br>Allow one sign error.              |
|          | $p = 1$ and $p = 6$                                                                                                                   | <b>A1</b>    |                                                                                            |
|          | For $p = 1$ , $d = -3$ , or $p = 6$ , $d = 12$                                                                                        | <b>DM1</b>   | For finding $d$ for one progression.                                                       |
|          | $a = 7 - (-3) \times 9 [= 34]$ $a = 12 - 12 \times 9 [= -96]$                                                                         | <b>DM1</b>   | For finding $a$ for one progression.                                                       |
|          | $u_{20} = 34 - 3 \times 19 = -23$ $u_{20} = -96 + 19 \times 12 = 132$                                                                 | <b>A1</b>    | For both.                                                                                  |
|          | Difference = 155                                                                                                                      | <b>A1)</b>   | Allow $-155$ .                                                                             |
|          | <b>OR</b><br>Solution 2<br><b>Alternative method for 2nd and 3rd M1</b><br>Lists at least three consecutive terms with position clear | <b>(DM1</b>  | E.g., 7, 4, 1 are 10th, 11th, 12th terms<br>or 34, 31, 28 are 1st, 2nd, 3rd terms.         |
|          | Lists at least three consecutive terms with position clear                                                                            | <b>DM1)</b>  | E.g. 12, 24, 36 are 10th, 11th, 12th terms,<br>or $-96, -84, -72$ are 1st, 2nd, 3rd terms. |
|          | Available marks                                                                                                                       | <b>6</b>     |                                                                                            |

## PUBLISHED

| Question | Answer                                                                                                                                                   | Marks      | Guidance                                                                                  |
|----------|----------------------------------------------------------------------------------------------------------------------------------------------------------|------------|-------------------------------------------------------------------------------------------|
| 7(a)     | $(r\sqrt{7})^2 = r^2 + (2r)^2 - 2 \times 2r \times r \times \cos \alpha \left[ \Rightarrow \alpha = \cos^{-1} \frac{4+1-7}{2 \times 2 \times 1} \right]$ | <b>M1</b>  | Correct use of cosine rule.                                                               |
|          | $= \frac{2}{3}\pi$ or $120^\circ$                                                                                                                        | <b>A1</b>  |                                                                                           |
|          | Arc length $= 2r \times \frac{2}{3}\pi$ or $\frac{120}{360} \times 2\pi \times 2r$                                                                       | <b>M1</b>  | Use of $r\theta$ with <i>their</i> angle.<br>Using <i>their</i> numeric angle in radians. |
|          | Total perimeter $= \left(\frac{4}{3}\pi + 1 + \sqrt{7}\right)r$                                                                                          | <b>A1</b>  | Or exact equivalent.<br>Allow unsimplified.                                               |
|          |                                                                                                                                                          | <b>4</b>   |                                                                                           |
| 7(b)     | Sector area $= \frac{1}{2} \times (2r)^2 \times \frac{2}{3}\pi$ or $\frac{120}{360} \times \pi \times (2r)^2$                                            | <b>M1*</b> | Using <i>their</i> numeric angle.                                                         |
|          | Triangle area $= \frac{1}{2} \times r \times 2r \times \sin \frac{2}{3}\pi$ or $= \frac{1}{2} \times r \times 2r \times \sin 120$                        | <b>M1*</b> | Using <i>their</i> numeric angle.                                                         |
|          | Area of shaded section $= \frac{1}{2} \times (2r)^2 \times \frac{2}{3}\pi - \frac{1}{2} \times r \times 2r \times \sin \frac{2}{3}\pi$                   | <b>DM1</b> | For subtraction of <i>their</i> areas.                                                    |
|          | $= \left(\frac{4}{3}\pi - \frac{\sqrt{3}}{2}\right)r^2$                                                                                                  | <b>A1</b>  | Or exact equivalent.<br>Allow unsimplified with $\sin \frac{2}{3}\pi$ evaluated.          |
|          |                                                                                                                                                          | <b>4</b>   |                                                                                           |

## PUBLISHED

| Question | Answer                                                                                                                                  | Marks      | Guidance                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|----------|-----------------------------------------------------------------------------------------------------------------------------------------|------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 8        | Meet when $\frac{5}{2x-1} = 7-2x \Rightarrow -4x^2 + 16x - 12 = [0] [\Rightarrow x^2 - 4x + 3 = 0]$                                     | <b>M1</b>  | For expressing as a three-term quadratic.<br>All terms need not be on the same side.                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|          | $x=1$ and $x=3$                                                                                                                         | <b>A1</b>  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|          | Volume = $[\pi] \int \left( (7-2x)^2 - \left( \frac{5}{2x-1} \right)^2 \right) dx$                                                      | <b>M1*</b> | Subtracting two volumes.<br>Condone subtraction the wrong way round.                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|          | $[\pi] \left( \left\{ 49x - 14x^2 + \frac{4}{3}x^3 \right\} [-] \left\{ -\frac{1}{2} \right\} \left\{ \frac{25}{2x-1} \right\} \right)$ | <b>B1</b>  | Or $\left\{ -\frac{1}{6}(7-2x)^3 \right\}$ . Allow unsimplified.                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|          |                                                                                                                                         | <b>B1</b>  | Allow unsimplified.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|          |                                                                                                                                         | <b>B1</b>  | Allow unsimplified.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|          | $[\pi] \left( 57 + \frac{5}{2} \right) - [\pi] \left( \frac{109}{3} + \frac{25}{2} \right)$ Minimum acceptable                          | <b>DM1</b> | Use of <i>their</i> limits in an attempt at integration. Condone one sign error or missing brackets.<br>Note<br>$(49 \times 3 - 14 \times 3^2 + \frac{4}{3} \times 3^3 + \frac{1}{2} \left( \frac{25}{2 \times 3 - 1} \right)) - (49 \times 1 - 14 \times 1^2 + \frac{4}{3} \times 1^3 + \frac{1}{2} \left( \frac{25}{2 \times 1 - 1} \right))$ $= (147 - 126 + 36 + \frac{5}{2}) - (\frac{109}{3} + \frac{25}{2})$ $= (57 + \frac{5}{2}) - (\frac{109}{3} + \frac{25}{2})$ $= \frac{119}{2} - \frac{293}{6}$<br>Or $\frac{62}{3} - 10$ |

## PUBLISHED

| Question | Answer                     | Marks     | Guidance                                                   |
|----------|----------------------------|-----------|------------------------------------------------------------|
| 8        | Volume = $\frac{32}{3}\pi$ | <b>B1</b> | Correct limits must be clear and integration correct, WWW. |
|          |                            | <b>8</b>  |                                                            |

| Question | Answer                                                      | Marks     | Guidance                                                                                                                                      |
|----------|-------------------------------------------------------------|-----------|-----------------------------------------------------------------------------------------------------------------------------------------------|
| 9(a)     | $y^3 = x \pm 8a$ or $x^3 = y \pm 8a$                        | <b>M1</b> |                                                                                                                                               |
|          | $[g^{-1}(x)] = \sqrt[3]{x+8a}$                              | <b>A1</b> | Do not allow “ $x = \dots$ ”.                                                                                                                 |
|          |                                                             | <b>2</b>  |                                                                                                                                               |
| 9(b)(i)  | $f(-1) = -4$                                                | <b>B1</b> |                                                                                                                                               |
|          | $g(-4) = (-4)^3 - 8a \Rightarrow (-4)^3 - 8a + 20a^2 [= 0]$ | <b>M1</b> | For $g(\text{their 'f(-1)'})$ and forming quadratic equation with all terms on one side.                                                      |
|          | $4(5a+8)(a-2)$                                              | <b>M1</b> | Solving <i>their</i> quadratic using suitable method, e.g. factorisation, quadratic formula, completing the square. M0 if roots are not real. |
|          | $a = 2$                                                     | <b>A1</b> | <b>SC B1</b> if M1M0.<br>Ignore answer of $a = -\frac{8}{5}$ .                                                                                |
|          |                                                             | <b>4</b>  |                                                                                                                                               |

## PUBLISHED

| Question | Answer                                                                             | Marks       | Guidance                                                                                                                                        |
|----------|------------------------------------------------------------------------------------|-------------|-------------------------------------------------------------------------------------------------------------------------------------------------|
| 9(b)(ii) | For either $gf(x) = (x-3)^3 - 16$ or $fg(x) = (x^3 - 16) - 3$                      | <b>B1FT</b> | In correct part of equation with $a > 0$ .<br>May be implied by correct equation.<br>Allow $gf(x) = (x-3)^3 - 8a$ or $fg(x) = (x^3 - 8a) - 3$ . |
|          | $(x-3)^3 - 16 = (x^3 - 16) - 3 + k$                                                | <b>M1*</b>  | For forming equation with $a$ substituted with $a$ real or $8a$ eliminated.<br>gf and fg must be the correct way round                          |
|          | $x^3 - 9x^2 + 27x - 27 - 16 = x^3 - 19 + k \Rightarrow -9x^2 + 27x - 24 - k [= 0]$ | <b>DM1</b>  | For multiplying out and simplifying.<br>All terms to one side.<br>Allow one sign error only.                                                    |
|          | $27^2 - 4 \times 9 \times (24 + k) [< 0]$                                          | <b>M1</b>   | For use of $b^2 - 4ac$ correct for <i>their</i> $a, b, c$ from a three-term quadratic and including $k$ .                                       |
|          | $k > -3.75$ or $k > -\frac{15}{4}$ or $k > -\frac{135}{36}$                        | <b>A1</b>   | OE                                                                                                                                              |
|          |                                                                                    | <b>5</b>    |                                                                                                                                                 |

| Question | Answer                                                                    | Marks     | Guidance |
|----------|---------------------------------------------------------------------------|-----------|----------|
| 10(a)    | At $x = 2$ , $\frac{d^2y}{dx^2} = 12 - 6$ [ $> 0$ ] $\Rightarrow$ minimum | <b>B1</b> |          |
|          |                                                                           | <b>1</b>  |          |

## PUBLISHED

| Question  | Answer                                                                                                                         | Marks      | Guidance                                                                                 |
|-----------|--------------------------------------------------------------------------------------------------------------------------------|------------|------------------------------------------------------------------------------------------|
| 10(b)     | $\frac{dy}{dx} = x^3 - 6x + c$                                                                                                 | <b>B1</b>  | $\frac{dy}{dx}$ can be implied by later work. Allow $f'(x)$ or $y'$ .                    |
|           | $0 = 8 - 12 + c \Rightarrow c = 4 \Rightarrow \frac{dy}{dx} = x^3 - 6x + 4$                                                    | <b>M1</b>  | For substituting $\frac{dy}{dx} = 0$ at $x = 2$ .                                        |
|           | $y = \frac{1}{4}x^4 - 3x^2 + 4x + k$                                                                                           | <b>A1</b>  | Can be implied by later work.                                                            |
|           | $3 = 4 - 12 + 8 + k \Rightarrow k = 3$                                                                                         | <b>M1</b>  | For substituting (2, 3) into <i>their</i> “ $y = \dots$ ”                                |
|           | $y = \frac{1}{4}x^4 - 3x^2 + 4x + 3$                                                                                           | <b>A1</b>  | Must be an equation unless $y =$ seen earlier in the question. Allow “ $f(x) = \dots$ ”. |
|           |                                                                                                                                | <b>5</b>   |                                                                                          |
| 10(c)(i)  | $x^3 - 6x + 4 = (x - 2)(x^2 + 2x - 2) [= 0]$                                                                                   | <b>M1*</b> | For using <i>their</i> $k$ in the given relationship or factorising otherwise.           |
|           |                                                                                                                                | <b>DM1</b> | Solving quadratic using suitable method, e.g. quadratic formula.                         |
|           | $x = -1 \pm \sqrt{3}$                                                                                                          | <b>A1</b>  | <b>SC B1</b> if M1 M0.                                                                   |
|           |                                                                                                                                | <b>3</b>   |                                                                                          |
| 10(c)(ii) | At $x = -1 + \sqrt{3} \quad \left[ \frac{d^2y}{dx^2} \right] = 12 - 6\sqrt{3} - 6$ or $-4.39\dots [ < 0 ] \Rightarrow$ maximum | <b>B1</b>  | Must see $\frac{d^2y}{dx^2}$ evaluated.                                                  |
|           | At $x = -1 - \sqrt{3} \quad \left[ \frac{d^2y}{dx^2} \right] = 6\sqrt{3} + 12 - 6$ or $16.4\dots [ > 0 ] \Rightarrow$ minimum  | <b>B1</b>  | Must see $\frac{d^2y}{dx^2}$ evaluated.                                                  |
|           |                                                                                                                                | <b>2</b>   |                                                                                          |

## PUBLISHED

| Question | Answer                                                                                                                                                                                        | Marks        | Guidance                                                                                                           |
|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|--------------------------------------------------------------------------------------------------------------------|
| 11       | <b>EITHER</b><br>Solution 1<br>Gradients of normal lines are $\frac{16-10}{6-8} [= -3]$ and $\frac{8-10}{14-8} [= -\frac{1}{3}]$                                                              | <b>(M1)*</b> | For either normal gradient.                                                                                        |
|          | Gradients of tangent lines are $\frac{1}{3}$ and 3                                                                                                                                            | <b>DM1</b>   | For either tangent gradient. Must be clear that they are finding tangent gradient.                                 |
|          | Equation of tangent at (6, 16) is $y - 16 = \frac{1}{3}(x - 6) [y = \frac{1}{3}x + 14]$                                                                                                       | <b>A1</b>    |                                                                                                                    |
|          | Equation of tangent at (14, 8) is $y - 8 = 3(x - 14) [y = 3x - 34]$                                                                                                                           | <b>A1</b>    |                                                                                                                    |
|          | Meet when $3x - 34 = \frac{1}{3}x + 14 \Rightarrow 8x = 144 [\Rightarrow x = 18]$                                                                                                             | <b>M1</b>    | Equating <i>their</i> two tangents.                                                                                |
|          | $x = 18$                                                                                                                                                                                      | <b>A1</b>    |                                                                                                                    |
|          | $y = 3x - 34$ meets $x$ -axis at $\frac{34}{3}$ or $y = \frac{1}{3}x + 14$ meets $y$ -axis at 14                                                                                              | <b>B1</b>    | For either. Coordinates of $P$ or $R$ must be clear or indication of where the lines cross the $x$ - or $y$ -axes. |
|          | Finds $y$ -coordinate of $Q$ and uses Area = rectangle – triangles<br>$= 18 \times 20 - \frac{1}{2} \times 18 \times (20 - 14) - \frac{1}{2} \times \left(18 - \frac{34}{3}\right) \times 20$ | <b>M1</b>    | For correct attempt at area, e.g. trapezium – triangle, integration.                                               |
|          | $= 360 - 54 - \frac{200}{3} = \frac{718}{3}$ or $239\frac{1}{3}$                                                                                                                              | <b>A1)</b>   |                                                                                                                    |

## PUBLISHED

| Question | Answer                                                                                                                                                                                                                                        | Marks | Guidance |
|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|----------|
| 11       | <b>OR</b><br>Solution 2<br><b>Alternative method for M1 M1</b><br>$\frac{dy}{dx} = \frac{16-2x}{2y-20}$                                                                                                                                       | (M1*) |          |
|          | $\frac{16-2 \times 6}{2 \times 16-20} \left[ = \frac{1}{3} \right] \text{ or } \frac{16-2 \times 14}{2 \times 8-20} [= 3]$                                                                                                                    | DM1)  |          |
|          | <b>OR</b><br>Solution 3<br><b>Alternative method 1 for final M1 A1</b><br>Finds y-coordinate of Q and uses Area = trapezium – triangle<br>$= \left( \frac{14+20}{2} \right) \times 20 - \frac{\left( 18 - \frac{34}{3} \right) \times 20}{2}$ | (M1   |          |
|          | $= 306 - \frac{200}{3} = \frac{718}{3} \text{ or } 239\frac{1}{3}$                                                                                                                                                                            | A1)   |          |
|          | <b>OR</b><br>Solution 2<br><b>Alternative method 2 for final M1 A1</b><br>$\int_0^{18} \left( \frac{1}{3}x + 14 \right) dx - \int_{\frac{34}{3}}^{18} (3x - 34) dx \Rightarrow \frac{1}{6}x^2 + 14x - \left( \frac{3}{2}x^2 - 34x \right)$    | (M1   |          |
|          | $306 - \frac{200}{3} = \frac{718}{3} \text{ or } 239\frac{1}{3}$                                                                                                                                                                              | A1)   |          |

**PUBLISHED**

| Question | Answer                                                                                                                       | Marks       | Guidance                      |
|----------|------------------------------------------------------------------------------------------------------------------------------|-------------|-------------------------------|
| 11       | <b>OR</b><br>Solution 2<br><b>Alternative method 1 for final M1 M1 A1 A1</b><br>$(x - 8)(x_1 - 8) + (y - 10)(y_1 - 10) = 40$ | <b>(M1)</b> | May be implied by later work. |
|          | Substitutes (6, 16) or (14, 8)                                                                                               | <b>DM1</b>  |                               |
|          | $-2(x - 8) + 6(y - 10) = 40 \Rightarrow y = \frac{1}{3}x + 14$                                                               | <b>A1</b>   |                               |
|          | $6(x - 8) - 2(y - 10) = 40 \Rightarrow y = 3x - 34$                                                                          | <b>A1)</b>  |                               |
|          | Available marks                                                                                                              | <b>9</b>    |                               |