



Mark Scheme (Results)

Summer 2025

Pearson Edexcel International GCSE  
In Chemistry (4CH1) Paper 2CR

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2025

Question Paper Log Number P75823A

Publications Code 4CH1\_2CR\_2506\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2025

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a) (i)	helium and neon	both are required for the mark	1
(ii)	oxygen	ACCEPT O <sub>2</sub>	1
(iii)	carbon dioxide	ACCEPT CO <sub>2</sub>	1
(iv)	carbon monoxide	ACCEPT CO	1
(b)	M1 (bubble the gas through) limewater M2 which turns (from colourless to) cloudy/milky	M2 dependent on M1 ALLOW white precipitate	2
			Total = 6

Question number	Answer	Notes	Marks
2 (a)	(fractional) distillation		1
(b)	(surfacing) roads / road tar / tarmac/ roofs	ALLOW 'making roads'	1
(c) (i)	$C_4H_{10}$		1
(ii)	butane		1
(iii)	$C_nH_{2n+2}$		1
			Total = 5

Question number	Answer	Notes	Marks
3 (a) (i)	(a solution that) has as much solute/solid/substance as possible dissolved (at a particular temperature) OWTTE	ALLOW No more solute will dissolve	1
(ii)	the solubility decreases (as the temperature decreases)		1
(b)	M1 (mass of potassium permanganate) 1.25 (g)  M2 (solubility of potassium manganate) 5 (g/100g)	ALLOW M1×4  5 (g/100g) no working scores 2 marks	2
(c) (i)	glowing splint/spill relights	REJECT burning/lit splint	1
(ii)	197	IGNORE any units given	1
(d)	any one from:  electron acceptor/gains electrons  oxygen donor/loses oxygen	IGNORE 'oxidises another substance whilst being reduced itself'	1
			Total = 7

Question number	Answer	Notes	Marks
4 (a) (i)	$2\text{Li(s)} + 2\text{H}_2\text{O(l)} \rightarrow 2\text{LiOH(aq)} + \text{H}_2\text{(g)}$	All four correct scores 2 2 or 3 correct scores 1	2
(ii)	any two from:  M1 effervescence  M2 gets smaller  M3 floats  M4 moves	ALLOW fizzing / bubbling IGNORE hydrogen/gas produced  ALLOW disappears / dissolves  moves on the surface scores <b>M3</b> and <b>M4</b>  REJECT melts/molten sphere REJECT flame/burning	2
(iii)	M1 (amount of hydrogen) = $375 \div 24000$ OR 0.015625 (mol)  M2 (amount of lithium) = 0.03125 (mol)  M3 (mass of lithium) 0.21875 (g)	ALLOW $M1 \times 2$  ALLOW $M2 \times 7$  ALLOW any number of significant figures apart from 1 REJECT incorrect rounding  ALLOW any answer that rounds to 0.22g  Answer with no working scores M3 only	3



<p>(b)</p>	<p>(i) any one from:</p> <p>potassium melts / turns into a ball / potassium moves faster / faster effervescence / (lilac/purple/pink) flame</p> <p>(ii) M1 potassium has a bigger atom / potassium has more shells / outer electron is further from the nucleus</p> <p>M2 the <u>outer</u> (shell) electron is less attracted to the nucleus <b>OR</b> the <u>outer</u> (shell) electron is more shielded (from the nuclear attraction)</p> <p>M3 (so outer) electron (in potassium) more easily lost</p>	<p>ALLOW disappears/dissolves faster</p> <p>REJECT more outer shells for M1</p> <p>Should be comparative for M3</p> <p>IGNORE electrons (plural) in M1 and M2 but do not allow electrons in M3</p> <p>ALLOW reverse argument throughout for lithium</p>	<p>1</p> <p>3</p>
			<p>Total = 11</p>

Question number	Answer	Notes	Marks
5 (a)	M1 electrolysis  M2 as sodium is more reactive than carbon	REJECT electrolysis of an aqueous solution  ALLOW it is above carbon in the reactivity series  ALLOW carbon cannot displace sodium  M2 dependent on M1	2
(b)	M1 layers / rows (of atoms, ions or particles)  M2 can slide over one another	REJECT if mention of molecules or intermolecular forces for M1  REJECT electrons sliding for M2	2
(c)	M1 sodium loses electrons and oxygen gains electrons  M2 two atoms of sodium lose one electron and oxygen gains 2 electrons	ACCEPT correctly labelled diagrams  Mention of sharing electrons scores 0	2
(d)	Sodium:  M1 (giant) metallic structure/lattice  M2 (electrostatic) attraction between positive ions and delocalised electrons  Sodium oxide:  M3 giant ionic structure OR (giant) ionic lattice  M4 (electrostatic) attraction between oppositely charged ions  Comparison:  M5 more energy is required to overcome the bonds in sodium oxide than sodium ORA	REJECT Intermolecular forces for M1 and M2  REJECT Intermolecular forces/covalent/metallic for M3 and M4  M5 is comparative mark that is independent of M1-M4	5
			Total = 11

Question number	Answer	Notes	Marks
6 (a)	<p>any two of the following pairs of points:</p> <p><b>M1</b> use a pipette (instead of a measuring cylinder)</p> <p><b>M2</b> so the volume of alkali added is more precise</p> <p>OR</p> <p><b>M1</b> add the acid dropwise towards the endpoint</p> <p><b>M2</b> so the exact endpoint is seen</p> <p>OR</p> <p><b>M1</b> swirl the flask</p> <p><b>M2</b> so the acid and alkali are fully mixed/reacted</p> <p>OR</p> <p><b>M1</b> repeat the titration</p> <p><b>M2</b> so concordant results are obtained OR so anomalous results can be ignored</p> <p>OR</p> <p><b>M1</b> do the titration on a white tile</p> <p><b>M2</b> so the colour change can easily be seen</p> <p>OR</p> <p><b>M1</b> rinse the burette with acid</p> <p><b>M2</b> so that the concentration of acid in the burette is not affected</p>	<p>IGNORE choice of indicator</p> <p>M2 is dependent on M1</p> <p>ALLOW accurate</p> <p>ALLOW stir</p>	4
(b)	<p><b>M1</b> (amount of potassium hydroxide)  <math>(25 \times 0.150) \div 1000</math> OR 0.00375 (mol)</p> <p><b>M2</b> <math>0.00375 \div 3</math> OR 0.00125 (mol)</p> <p><b>M3</b> <math>(0.00125 \div 0.04) \times 1000</math></p> <p><b>M4</b> 31.3(cm<sup>3</sup>)</p>	<p>ALLOW M1÷3</p> <p>ALLOW ecf from M2</p> <p>M3 to 3 sig figs</p> <p>31.3 (cm<sup>3</sup>) scores 4  31.25 (cm<sup>3</sup>) scores 3  93.8 (cm<sup>3</sup>) scores 3  93.75 (cm<sup>3</sup>) scores 2</p>	4
(c) (i)	$2K_3PO_4 + 3CaCl_2 \rightarrow Ca_3(PO_4)_2 + 6KCl$		1

<p>(ii)</p>	<p>M1 (amount of calcium phosphate) 0.0200 (mol)</p> <p>M2 (mass of calcium phosphate) 6.2 (g)</p> <p>M3 <math>(5.7 \div 6.2 \times 100) = 91.9 \%</math></p> <p>OR</p> <p>M1 (amount of calcium phosphate) = <math>5.7/310</math> OR 0.0184 (mol)</p> <p>M2 <math>0.0184 \times 2</math> OR 0.0368 (mol)</p> <p>M3 <math>(0.0368/0.04 \times 100) = 91.9 \%</math></p>	<p>ALLOW <math>M1 \times 310</math></p> <p>ALLOW any number of significant figures from 2</p> <p>ALLOW <math>M1 \times 2</math></p> <p>ALLOW any number of significant figures from 2</p>	<p>3</p>
<p>(iii)</p>	<p>M1 filter (the calcium phosphate)</p> <p>M2 wash the residue (with distilled water)</p> <p>M3 leave to dry (on a windowsill) / dry between filter papers / dry in a desiccator / dry in a warm oven</p>	<p>If evaporation to form crystals/a soluble salt method then <b>M3 only</b> for a suitable drying method.</p> <p>REJECT hot oven or direct heating with Bunsen Burner for M3</p> <p>REJECT if solid is washed again after drying for M3</p>	<p>3</p>
			<p>Total = 15</p>

Question number	Answer	Notes	Marks	
7	(a) (i)	yeast	ALLOW zymase/enzymes from yeast	1
	(ii)	C 30°C  A is not the correct answer because 0°C is too low for fermentation B is not the correct answer because 10°C is too low for fermentation D is not the correct answer because 100°C is too high for fermentation		1
	(iii)	An explanation using either of the following linked pairs:  M1 oxygen (in the air) would react with ethanol/ alcohol  M2 to form ethanoic acid  <b>OR</b>  M1 (in the presence of air) the glucose would react with the oxygen  M2 so ethanol does not form	ALLOW ethanol would be oxidised  ALLOW to form carboxylic acid ALLOW to form vinegar/acetic acid  ALLOW correct reference to aerobic respiration  ALLOW forms CO <sub>2</sub> and H <sub>2</sub> O	2
(b) (i)	M1 $346 + (5 \times 414) + 358 + 463 + (3 \times 498)$ <b>OR</b> 4731 M2 $(4 \times 804) + (6 \times 463)$ <b>OR</b> 5994 M3 -1263 (kJ/mol)  <b>OR</b> M1 $346 + (5 \times 414) + 358 + (3 \times 498)$ <b>OR</b> 4268 M2 $(4 \times 804) + (5 \times 463)$ <b>OR</b> 5531 M3 -1263 (kJ/mol)	-1263 kJ/mol scores 3 ALLOW ecf for M3	3	

<p>(ii)</p>	 <p>M1 right-hand line below left-hand line</p> <p>M2 correct names/balanced formulae of both reactants</p> <p>M3 correct names/balanced formulae of both products</p>	<p>IGNORE any activation energy hump</p> <p>IGNORE enthalpy change arrow</p> <p>If only use words reactants (on left) and products (on right) award 1 mark from M2 and M3</p> <p>If no/incorrect coefficients then 1 mark from M2 and M3</p>	<p>3</p>
<p>(iii)</p>	<p>An explanation that links together the following two points:</p> <p>M1 more energy is given out when the bonds are made</p> <p>M2 than is taken in when the bonds are broken ACCEPT correct reverse argument</p> <p><b>OR</b></p> <p>M1 breaking bonds is endothermic/takes in energy AND making bonds is exothermic/releases energy</p> <p>M2 the energy released is more than the energy taken in</p>	<p>If state/imply that energy required to make bonds OR If state/imply that energy released when bonds are broken scores 0/2</p> <p>M2 dep on M1</p>	<p>2</p>
<p>(c) (i)</p>	<p>M1 displayed formula for ethanoic acid</p>  <p>M2 displayed formula for water/H<sub>2</sub>O</p>	<p>ACCEPT any shape</p>	<p>2</p>

(ii)	ethyl ethanoate	IGNORE hyphens	1
			Total = 15

